

Read the label before opening the container.

For Full particulars, see enclosed leaflet.

StarBug

South Africa Reg.No: L 8008 Act No. 36 of 1947

A systemic suspension concentrate insecticide for the control of aphids, red scale, thrips, leafminers and psylla on citrus trees; aphids on potatoes, cucurbits, cruciferae and roses; woolly aphids on apples and whitefly on tomatoes.

IRAC INSECTICIDE GROUP CODE: 4A

ACTIVE INGREDIENT:

Imidacloprid (neonicotinoid) 350 g/l

Product Information Tel no: 082 801 6759
In case of poisoning: 082 446 8946

www.enviro-crop.co.za

enviro
efficiency • quality • competitiveness

Batch No:
Date of Manufacture:

UN No. 2902

Registration holder:

Erintrade cc t/a RT Chemicals
CK 2001/036403/23
10 Ninth Avenue, Industria, Kroonstad, 9500
Tel: 056 213 1967

06082012-07/12



**HARMFUL
SKADELIK**



WARNINGS:

- Allow the following interval between the last **StarBug** application and harvest:

Apples: Foliar Application	70 days	Citrus:	212 days
Broccoli:	76 days	Cucurbits:	100 days
Brussels Sprouts:	91 days between planting and harvest	Potatoes:	120 days
Cabbage:	91 days between planting and harvest	Tomatoes:	23 days
Cauliflower:	136 days between planting and harvest		

NOTE: The recommended withholding period between last application and harvest meets local residue requirements but may not meet export residue requirements. If the treated crop is intended for the export market, consult the relevant exporting body regarding pre-harvest interval requirements before the application of this product,

- Handle with care.
- Poisonous when absorbed through the skin or swallowed or inhaled.
- Toxic to fish and wildlife.
- Store in a cool place away from food and feedstuffs.
- Keep out of reach of children, uninformed persons and animals.
- **In case of poisoning call a doctor and make label available to him/her.**
- **RE-ENTRY:** Do not enter treated area within 1 day after treatment unless wearing protective clothing.
- **AERIAL APPLICATION:** Notify all inhabitants of the immediate area to be sprayed and issue the necessary warnings. Do not spray over or allow drift to contaminate adjacent areas or water and rivers. Please refer to the SANS Code of Practice for the aerial application of agricultural chemicals.

Although this insecticide has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions; quality of dilution/application water; compatibility with other substances not indicated on the label and the occurrence of resistance of the pest against the insecticide concerned as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the insecticide concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

PRECAUTIONS:

- Wear protective clothing (face mask, overall, gloves, boots) when mixing and applying this product.
- Do not inhale fumes.
- Wash contaminated clothing daily.
- Wash with soap and cold water after use or accidental skin contact.
- Do not eat, drink or smoke whilst mixing, applying or before washing hands and face.
- Clean applicator after use, dispose of wash water where it will not contaminate crops, grazing, rivers and dams.
- Rinse empty container three times with a volume of water equal to at least one tenth of that of the container and add the rinsing water to the contents of the tank before disposing of the container in the prescribed manner.
- Destroy empty container by perforation and flattening and never use for any other purpose.
- Prevent contamination of food, feedstuffs, drinking water and eating utensils.

RESISTANCE WARNING:

For resistance management, **StarBug** is a group code 4A insecticide. Any insect population may contain individuals naturally resistant to **StarBug** and other group code 4A insecticides. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly and consecutively. These resistant insects may not be controlled by

StarBug or any other group code 4A insecticides.

To delay insecticide resistance:

- Avoid exclusive repeated and consecutive use of insecticides from the same insecticide group code. Alternate with products from different insecticide group codes.
 - Do not apply **StarBug** or any product that belongs to the same chemical group, more than once per year on citrus (bearing, non-bearing and nursery trees).
 - Integrate other control methods (chemical, cultural, biological) into insect control programs.
- For specific information on resistance management contact the registration holder of this product.

SYMPTOMS OF HUMAN POISONING:

Apathetic state, depression, muscular tone, respiratory disturbances and trembling and muscular cramps in severe cases of poisoning.

FIRST AID TREATMENT:

Remove the patient from source of poisoning to a well ventilated area and keep him/her quiet and reassured.

Remove contaminated clothing. Rinse the contaminated body area thoroughly with plenty of soap and cold water. Do not rub skin.

Flush contamination out of eyes with clean water for at least 15 minutes.

If swallowed, induce vomiting by tickling the back of the throat/uvula. Repeat until vomit fluid is clear and free from the smell of the insecticide. Administer artificial respiration or closed chest cardiac massage if necessary. Do not apply direct mouth to mouth respiration. Never give anything by mouth to an unconscious person. **Take the patient to the nearest physician immediately.**

NOTE TO PHYSICIAN:

Treat symptomatically. Monitor blood pressure and pulse rate frequently (bradycardia and hypotonia are possible). Provide supportive measures for respiratory and cardiac action. Give artificial respiration if signs of paralysis appear. Additional measures involve the elimination of the substance from the body or acceleration of its excretion (gastro lavage, saline laxatives).

DIRECTIONS FOR USE: USE ONLY AS INDICATED.

Compatibility:

StarBug is compatible with Atador (Reg. No. L7167; Registration holder: Erintrade cc t/a RT Chemicals, fenamiphos 400 EC) when applied on potatoes.

Mixing Instructions:

Pour the required quantity of **StarBug** into the spray water and agitate continuously.

Do not allow the **StarBug** mixture to stand any length of time before using.

Do not use brack or hard water.

Aerial Application:

Aerial application may only be done by a registered aerial application operator using a correctly calibrated, registered aircraft according to the instructions of SANS Code 0118 (Aerial Application of Agricultural Remedies). It is important to ensure that the spray mixture is distributed evenly over the target area and that the loss of spray material during application is restricted to a minimum. It is therefore essential that the following criteria are met:

- Volume: A minimum volume of 30 ℓ spray mixture per hectare is recommended. As this product has not been evaluated at a reduced volume rate, the registration holder cannot guarantee efficacy, or be held responsible for any adverse effects if this product is applied aerially at a lower volume rate than recommended above.
- A droplet coverage of 30 to 40 droplets per cm² must be recovered on the target.

- A droplet spectrum with a VMD of 250 to 280 microns is recommended. Ensure that the production of fine droplets (less than 150 microns) is restricted to a minimum.
- The height of the spray boom should be maintained at 3 to 4 metres above the target. Do not spray when aircraft is in a climb, at the top or during a dive, or when banking.
- Use suitable atomising equipment (hydraulic nozzles or rotary atomisers) that will produce the desired droplet size and coverage, but which will ensure the minimum loss of product through drift. The operator must use a setup that will produce a droplet spectrum with the lowest possible relative span.
- All nozzles/atomisers should be positioned within the inner 60 to 75 % of the wingspan to prevent droplets from entering the wingtip vortices.
- Do not spray if the wind speed exceeds 15 km/h.
- The difference between the wet and dry bulb temperature readings as determined by a whirling hygrometer, must not exceed 8°C.
- Do not spray under turbulent, unstable conditions during the heat of the day when rising thermals and downdraughts occur.
- Spraying under temperature inversion conditions (spraying in or above the inversion layer) and/or high humidity conditions (relative humidity 80 % and above) may lead to the following:
 - reduced efficacy due to suspension and evaporation of small droplets in the air (inadequate coverage).
 - damage to other sensitive crops and/or non-target areas through drifting of the suspended spray cloud away from the target field.
- Ensure that fields are accurately marked and that the aerial spray operator knows exactly which fields to spray.
- Obtain an assurance from the aerial spray operator that the above requirements will be met and that a logbook record of the application will be made.

CROP / PEST:	DOSAGE RATE:	REMARKS:
CUCURBITS Aphids (Various species)	Ground Application: 200 mℓ/100 ℓ water	Apply 30 mℓ of the mixture per planting hole during planting or immediately after planting. Additional sprays for aphid control may be necessary later in the season should aphids appear.
CRUCIFERAE Cabbage aphid (<i>Brevicoryne brassicae</i>)	Ground Application: 380 mℓ/100 ℓ water	Apply 600 mℓ of the StarBug suspension per polystyrene seedling tray as a drench over the plants not more than 24 hours prior to transplanting.

NOTES:

1. The above application rates are recommended per polystyrene seedling tray containing 200 seedlings. The seedling trays referred to are those with a standard outside dimension of 673 x 342 mm.
2. Application rates can be proportionally reduced for seedling trays containing less than 200 seedlings.
3. The mixture should be stirred thoroughly before and during application to ensure an even application.

CROP / PEST:	DOSAGE RATE:	REMARKS:
ROSES Aphids (<i>Macrosiphum rosae</i> and <i>Metopolophium porosum</i>)	Soil drench: 0,5 mℓ / plant (for 4 - 6 weeks control) 1,0 mℓ/plant (for season long control)	Apply as a soil drench in spring. Mix 10 mℓ StarBug with 10 litres water and apply 1 litre of this mixture on the soil around the base of the stem of each plant. Irrigate within 24 hours after application. Follow-up applications may be necessary, depending on the amount of rainfall or irrigation and soil type.

CROP / PEST:	DOSAGE RATE:	REMARKS:
APPLES Woolly aphid <i>(Eriosoma lanigerum)</i>	Soil Application: <u>Trees up to 6 years old:</u> 1 ml/tree	Apply in September or within 3 weeks after flowering.
	<u>Trees 7 to 15 years old:</u> 3 ml/tree	Apply within 3 weeks after flowering. Apply the StarBug in 1 litre water to the soil directly around the base of the tree trunk. The application area should be free of weeds, organic matter and mulch. Irrigate within 24 hours after application. Will also suppress woolly aphid on the roots.
	Foliar Spray: 60 ml/100 l water (1,5 - 2,1 l/ha in 2500 - 3500 l water/ha)	Apply StarBug as a full cover high volume foliar spray during approximately mid-December. StarBug may be applied as a foliar spray to Top Red, Granny Smith, Royal Gala and Golden Delicious apples.

NOTES:

- Do not apply to trees older than 15 years.
- Trees should be healthy, stress free and regularly irrigated before and after application. Any factor that will stress the tree will adversely affect the uptake and distribution of the active ingredient in the tree.
- Do not apply the soil drench treatment through the irrigation system.

CROP / PEST:	DOSAGE RATE:	REMARKS:
POTATOES Aphid <i>(Myzus persicae)</i>	Ground Application: 4,5 ml/100 m row length	Spray the StarBug mixture to the soil surface in the planting furrow from ridge-tip to ridge-tip <u>before</u> planting of the tubers. Use an 8006 or 8008 flat fan nozzle and apply in 3 to 4 liters water per 100 metre furrow length. The application may also be made after planting but before emergence over the row. Irrigate immediately afterwards with \pm 15 mm water. StarBug may also be used in a tank mixture with Atador for the control of aphids and nematodes.
	3,0 ml/100 m row length	When StarBug is mixed with Atador , the StarBug dosage can be reduced to 3,0 ml per 100 metres row length. Warning: Do not apply StarBug mixtures directly to the tubers. Apply in the open furrow before planting of the tubers.
TOMATOES Whitefly nymph <i>(Trialeurodes vaporariorum)</i> For use on tomatoes in enclosed areas only, e.g. glass houses, tunnels, etc.	Soil drench: 0,05 ml/plant	Apply as a soil drench, 1 day after transplanting. Mix 10 ml StarBug with 20 litres water and apply 100 ml of this mixture to the soil around the base of the stem of each plant. Irrigate within 24 hours after application. Repeat application 28 days later.
		Apply as a soil drench in spring or at the first signs of infestation. Mix 5 ml StarBug with 10 litres water and apply 1 litre of this mixture on the soil around the base of the stem of each plant. Irrigate within 24 hours after application.

CROP / PEST:	DOSAGE RATE:	REMARKS:
<p>CITRUS</p> <p>Bearing Trees not older than 15 years (excluding Kumquat)</p> <p>Aphid (<i>Toxoptera citricidus</i>), Red scale (<i>Aonidiella aurantii</i>) & Psylla (<i>Trioza erythrae</i>)</p> <hr/> <p>Aphid (<i>Toxoptera citricidus</i>)</p>	<p>Soil drench:</p> <p>9 mL/tree</p> <hr/> <p>6 mL/tree</p>	<p>Apply during August/September at green tip to white bud growth stage of the trees. Dilute the required volume in 1 litre water and apply around the base of the trunk with a jug. Irrigate within 24 hours after application. When used on lemon trees, apply only after removal of all fruit (including immature fruit). In lemons control can only be expected on the first fruit set after application.</p>
<p>Non-bearing Trees</p> <p>Aphid (<i>Toxoptera citricidus</i>)</p> <hr/> <p>Leaf miner (<i>Phyllocnistis citrella</i>)</p>	<p>Soil drench:</p> <p>6 mL/tree</p> <hr/> <p>8 mL/tree</p>	<p>Apply during August/September in 1 litre water and apply around the base of the trunk with a jug. Irrigate within 24 hours after application.</p>
<p>Nursery Trees</p> <p>Aphid (<i>Toxoptera citricidus</i>) & Leaf miner (<i>Phyllocnistis citrella</i>)</p> <hr/> <p>Thrip (<i>Scirtothrips aurantii</i>)</p>	<p>Soil drench:</p> <p>1,5 mL/seedling</p> <hr/> <p>3,0 mL/seedling</p>	<p>Apply in 200 mL water 2 days after transplanting of the grafted root stock seedlings into plastic bags. Ensure that the StarBug application does not take place prior to grafting.</p>
<p>NOTES:</p> <ul style="list-style-type: none"> • Trees should be healthy, stress free and regularly irrigated before and after application. • Do not apply the soil drench treatment through the irrigation system. • Do not apply to trees immediately after pruning. 		