


# TERMITE STOP

Reg. No. L5437 Act 36/1947

**A ready to use bait with stomach action  
for the control of harvester termite,  
*Hodotermes mossambicus*.**

**Active Ingredient:** Sodium Fluorosilicate 50 g/kg



Registered and Formulated by:



(Pty) Ltd./ (Edms) Bpk.

Co. Reg. No.: 1961/001744/07  
PO Box 3131, Kenmare 1745  
Tel: 011 769 1300

**BAIT**



**CAUTION**



## **WARNINGS:**

Toxic when swallowed. Store in a cool place away from food and feedstuff. Keep out of reach of children, animals and uninformed persons.

## **PRECAUTIONS:**

Wash with soap and water after handling. Prevent contamination of food, feedstuffs, drinking water and eating utensils. Destroy empty container and do not use for any other purpose.

## **DIRECTIONS FOR USE:**

Use only as directed.

Time of treatment: Usually termites are active during Winter from 10h00 until sunset and during the Summer activity is usually after sunset.

The best period to scatter TERMITE STOP bait is during March to June, preferably when termites are active.

## **GENERAL INSTRUCTIONS:**

- Avoid wet conditions such as rain and irrigation until most bait is removed.
- Sharp smelling substances such as fuel, oil and some pesticides could contaminate the bait and repel termites.
- The smell on human hands is repellent. It is therefore advisable to wear latex gloves when handling the bait, or scatter the bait directly from the container.

## **DIRECTIONS FOR APPLICATION**

CROP	DOSAGE
Lawns	1g / foraging hole 250g / 125m <sup>2</sup>

Spread bait in small heaps (size of a matchbox) over and around the infested area, especially in close vicinity of foraging holes where termites are active. Termite activity declines after application with only white nymphs active, but they soon disappear. The area must be inspected 30 – 60 days after treatment and spots where termites still occur must be retreated.

Batch No.:

Date of Manufacture: